

diosmin + hesperidin
MPFF (Micronized Purified Flavonoid Fraction)

daflon[®] 1000
900mg / 100mg Film-coated Tablet
Vasoprotective

Description of the Product

Film-coated tablets.

The drug product is a salmon colored, oblong, film-coated tablet of 23.3 mm in length and 8.3 mm in width, scored on both faces and finished to a mass of 1378 mg.

The score line is only to facilitate breaking for ease of swallowing and not to divide into equal doses.

What is in the Medicine?

The active substances are:

Micronized purified flavonoid fraction 1000 mg
Corresponding to:
Diosmin 90 percent 900 mg
Flavonoids expressed as
hesperidin 10 percent 100 mg
Mean moisture 40 mg
For one film-coated tablet.

What is this Medicine used for?

This medicine is a venotonic (it increases venous tone) and a vasculoprotector (it increases resistance in small blood vessels).

This medicine is recommended for the treatment of venous circulation disorders (heavy legs, pain, restless legs) and functional symptoms related to acute hemorrhoidal attack.

If the hemorrhoid symptoms do not disappear within 15 days, talk to your doctor.

How much and how often should you use this Medicine?

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Venous insufficiency: 1 tablet daily, at mealtime.

Hemorrhoidal attack: The recommended dose is 3 tablets per day for 4 days and then 2 tablets per day for the next 3 days, at mealtimes.

Oral route.

When should you not take this Medicine?

If you are allergic (hypersensitive) to micronized purified flavonoid fraction or any of the other ingredients of this medicine.

Care that should be taken when taking this Medicine


Warnings and precautions

Talk to your doctor or pharmacist before taking this medicine.

The administration of this product for the symptomatic treatment of acute hemorrhoids does not preclude treatment for other anal conditions. The treatment must be short-term. If symptoms do not subside promptly, a proctological examination should be performed and the treatment should be reviewed.

Children

Not applicable.

GENERAL INFORMATIONS		SPOTS COLORS	NUMBER PAGES
PIL_FT_B_#02 160 x 210 mm (D1)_DAFLON_1000_LGS_005 14/10/2021 5915_05.01 PHILIPPINES (PHL)	Font size: 10/11 pts Font size of the section headings: 10 pts Font type: Helvetica Neue LT Pro Cnd	 PMS 287 U	 4 pages

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy: As a precautionary measure, it is preferable to avoid the use of this medicine during pregnancy.

Breast-feeding: In the absence of data in breast-feeding women, it is preferable to avoid breast-feeding while taking this treatment.

Sports

Not applicable.

Driving and using machines

Not applicable.

Undesirable Effects of this Medicine

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The frequency of the possible undesirable effects listed below is defined using the following system:

- Very common (affects more than 1 user out of 10)
- Common (affects 1 to 10 users out of 100)
- Uncommon (affects 1 to 10 users out of 1,000)
- Rare (affects 1 to 10 users out of 10,000)
- Very rare (affects less than 1 user out of 10,000)
- Frequency not known (cannot be estimated from the available data)

Nervous system disorders

Rare: dizziness, headaches, discomfort.

Gastrointestinal disorders

Common: diarrhea, dyspepsia (indigestion), nausea, vomiting.

Uncommon: colitis (inflammation of the large intestine).

Frequency not known: abdominal pain.

Skin and subcutaneous tissue disorders

Rare: rash, pruritus (itching), urticaria (hives).

Frequency not known: isolated face, eyelids and lips

edema (swelling). Exceptionally, Quincke's edema (swelling of the face, lips, mouth, tongue or throat which could cause breathing difficulties).

Reporting of Adverse Drug Reaction

For suspected adverse drug reaction, report to the FDA at www.fda.gov.ph

Seek medical attention immediately at the first sign of any adverse drug reaction.

What other medicine or food should be avoided while taking this Medicine?

Other medicines

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines obtained without prescription.

No clinically relevant drug interaction has been reported since the marketing of the product.

Food and drink

Not applicable.

What should you do if you miss a dose?

Not applicable.

Signs and Symptoms of Overdose

The experience of overdoses with this medicine is limited but reported symptoms include diarrhea, nausea, abdominal pain, pruritus and rash.

What to do when you have taken more than the recommended dosage?

Consult your doctor or pharmacist immediately.

Management of overdose should consist of treatment of clinical symptoms.

How should you keep this Medicine?

Store at temperatures not exceeding 30°C.

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the box. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask a pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

When should you consult your doctor?

Talk to your doctor or pharmacist before taking this medicine.

Les Laboratoires Servier – France



Manufactured by:

Les Laboratoires Servier Industrie

905 route de Saran
45520 Gidy – France

Imported by:

Servier Philippines, Inc.

Unit AD, 11F, 8 Rockwell
Hidalgo Drive, Rockwell Center
Brgy. Poblacion, Makati, Metro Manila

Distributed by:

Zuellig Pharma Corporation

Km. 14 West Service Road
South Super Highway cor. Edison Ave.
Sun Valley, Parañaque City

Reg. No.: DR-XY46549

Date of First Authorization: 8 April 2019
Date of Revision of Patient Information Leaflet:
24 November 2021

